

Technical note on seasonal adjustment for Index of industrial production (Consumer goods)

May 14, 2012

Contents

1	Index of industrial production (Consumer goods)	2
2	Steps in the seasonal adjustment procedure	2
2.1	Seasonal adjustment of IIP with X-12-ARIMA	4
2.2	Diagnostic checks	5
2.2.1	Validation of the automodel choice by X-12-ARIMA	5
2.2.2	Presence of identifiable seasonality	5
3	Sliding span diagnostics	6
4	Spectral representation	6
5	Accounting for India-specific moving holiday effects	7

List of Figures

1	IIP (Consumer goods) (Non seasonally adjusted)	2
2	Monthly growth rates across the years	3
3	IIP (NSA and SA)	4
4	ACF of residuals	5
5	IIP Spectral plot (NSA and SA)	7


List of Tables

1	Regression model for IIP (Consumer goods)	8
---	---	---

1 Index of industrial production (Consumer goods)

We analyse the monthly data for IIP (Consumer goods) from April, 1994 onwards. Figure 1 below shows the original plot of the series. The plot shows seasonal peaks. In a non-seasonally adjusted series, it is difficult to discern a trend as the seasonal variations may mask the important characteristics of a time series.

Figure 1 IIP (Consumer goods) (Non seasonally adjusted)




2 Steps in the seasonal adjustment procedure

Given that seasonality exists, it is important to model seasonality before the application of seasonal adjustment procedure. Seasonality in time series can be deterministic or stochastic. Stochastic seasonality can be stationary or non-stationary.

A visually appealing way of looking at the raw data is to plot the growth rates in each of the months across the years i.e the growth of April over March in each of the years from 1994 onwards. This gives us some idea of the presence of seasonal peaks, if any in the series. The nature of seasonality can also be inferred intuitively from the plot before the application of the testing procedures.

Figure 2 shows seasonal peaks in the month of March and December.

Figure 2 Monthly growth rates across the years



2.1 Seasonal adjustment of IIP with X-12-ARIMA

Seasonal adjustment is done with X-12-ARIMA method. Since the test results point towards deterministic seasonality, seasonal dummy is added in the regARIMA specification.

Figure 3 IIP (NSA and SA)





Figure 3 shows the non-seasonally and seasonally adjusted IIP (Consumer goods). The plot reveals that the seasonal peaks are dampened after seasonal adjustment.

2.2 Diagnostic checks

After seasonal adjustment, a series of diagnostic checks are performed through relevant tests and quality assessment statistics.

2.2.1 Validation of the automodel choice by X-12-ARIMA

Figure 4 ACF of residuals



A test of validation of the auto model choice by X-12-ARIMA is the randomness of the residuals of the fitted ARIMA model. The Ljung-Box test is conducted on the residuals of the fitted ARIMA model to check whether or not the residuals are white noise. The ACFs of the residuals are plotted to check for randomness.

Figure 4 does not reveal significant autocorrelation amongst the residuals.

2.2.2 Presence of identifiable seasonality

The statistic $M7$ shows the amount of moving seasonality present relative to stable seasonality. It shows the combined result for the test of stable and moving seasonality in the series. A value lesser than 1 is desirable to show identifiable seasonality in the series. The value for IIP (Consumer goods) is 0.4.

IIP (Consumer goods) series show identifiable seasonality on the basis of the $M7$ statistic.

3 Sliding span diagnostics

Sliding span diagnostics are descriptive statistics of how the seasonal adjustments and their month-to-month changes vary when the span of data used to calculate them is altered in a systematic way.

It is based on the idea that for a month common to more than one overlapping spans, the percent change of its adjusted value from the different spans should not exceed the threshold value and for a month common to more than one span, the difference between the month on month change from the different spans should not exceed the threshold value (the threshold value being 0.03).

Sliding span gives the percentage of months (A%) for which the seasonal adjustment is unstable (the difference in the seasonally adjusted values for a particular month from more than one span should not exceed 0.03). It also gives the percentage of months (MM%) for which the month on month changes of the seasonally adjusted values is unstable i.e exceeding the threshold value. The seasonal adjustment produced by the procedure chosen should not be used if $A\% > 25.0$ (> 15.0 is considered problematic) or if $MM\% > 40.0$.

Since the data span is short, sliding span diagnostic is not relied.


4 Spectral representation

Figure 5 shows the spectral plot of the growth rate of the unadjusted and seasonally adjusted series. Spectral plot, an important tool of the frequency domain analysis shows the portion of variance contributed by cycles of different frequencies.

The x-axis represent frequency from 0 to π (3.14). The seasonal frequencies are $\pi/6$ (0.52 on the x-axis), $\pi/3$ (1.04 on the x-axis), $\pi/2$ (1.57 on the x-axis), $2\pi/3$ (2.09 on the x-axis) and $5\pi/6$ (2.6 on the x-axis). In terms of periods (months); they are 12 months, 6 months, 4 months, 3 months and 2.4 months.

The figure at the lower panel shows that peaks at seasonal frequencies are eliminated after seasonal adjustment. For example the first peak at 0.52 correspond to 12 months which is eliminated after seasonal adjustment. Other peaks seen in the lower panel of the figure are not at seasonal frequencies.

Figure 5 IIP Spectral plot (NSA and SA)



5 Accounting for India-specific moving holiday effects

Accounting for moving holiday effect is a crucial component of pre-treatment of the series before the application of seasonal adjustment method. X-12-ARIMA is capable of handling the moving holiday effects through the inclusion of regressors for Easter Sunday, Labor Day, and Thanksgiving Day. These are important moving holidays for U.S time series.

We use the GENHOL program of X-12-ARIMA to analyse India-specific moving holiday effect. The program generates regressor matrices from holiday date file to enable X-12-ARIMA, estimation of complex moving holiday effects. It has the capability to generate regressors for before the holiday interval, surrounding the holiday interval and past the holiday interval.

The key assumption is that the fundamental structure of a time series changes for a fixed number of days before, after or for a fixed interval surrounding the holidays. We estimate the effect of Diwali which is an important moving holiday in Indian scenario. For estimating Diwali effect, we assume that the level of economic activity changes 5 days before Diwali (including the day on which Diwali falls). Regression variable for Diwali is found to be significant for IIP (Consumer goods).

Table 1 Regression model for IIP (Consumer goods)

Variable	Parameter estimate	Standard error	t-value
User defined			
Diwali	-0.0559	0.01367	-4.09

The results in Table 1 show significant trading day effect on IIP (Consumer goods) on account of Diwali. There is a significant drop in IIP (Consumer goods) around Diwali on account of loss of working days.