

Technical note on seasonal adjustment for Wholesale price index (Food articles)

May 14, 2012

Contents

1	WPI Food	2
1.1	Additive versus multiplicative seasonality	2
2	Steps in the seasonal adjustment procedure	2
2.1	Seasonal adjustment of WPI Food with X-12-ARIMA	3
2.2	Diagnostic checks	5
2.2.1	Validation of the automodel choice by X-12-ARIMA	5
2.2.2	Presence of identifiable seasonality	6
3	Year on year growth versus seasonally adjusted point on point growth	6
4	Spectral representation	7
5	Accounting for India-specific moving holiday effects	7

List of Figures

1	WPI of food articles (Non seasonally adusted)	2
2	Monthly growth rates across the years	3
3	WPI food articles (NSA and SA)	4
4	WPI (NSA and SA- point on point growth rates)	5
5	ACF of residuals	6
6	WPI (Food) Spectral plot (NSA and SA)	7


List of Tables

1	Year on year and point on point growth rates	9
---	--	---

1 WPI Food

We analyse the monthly data for Wholesale price index of food articles at the new base year from April, 2005 onwards. Figure 1 shows the original plot of the series. The plot shows seasonal peaks. In a non-seasonally adjusted series, it is difficult to discern a trend as the seasonal variations may mask the important characteristics of a time series.

Figure 1 WPI of food articles (Non seasonally adusted)



1.1 Additive versus multiplicative seasonality

X-12-ARIMA has the capability to determine the mode of the seasonal adjustment decomposition to be performed i.e whether multiplicative or additive seasonal adjustment decomposition is appropriate for the series. For the given series, multiplicative seasonal adjustment is considered appropriate on the basis of the model selection criteria.

2 Steps in the seasonal adjustment procedure

Given that seasonality exists, it is important to model seasonality before the application of seasonal adjustment procedure. Seasonality in time series can be deterministic or stochastic.

Stochastic seasonality can be stationary or non-stationary.

A visually appealing way of looking at the raw data is to plot the growth rates in each of the months across the years i.e the growth of April over March in each of the years from 1994 onwards. This gives us some idea of the presence of seasonal peaks, if any in the series. The nature of seasonality can also be inferred intuitively from the plot before the application of the testing procedures.

Figure 2 Monthly growth rates across the years




Figure 2 shows that the mean growth rate in April is higher than the other months.

2.1 Seasonal adjustment of WPI Food with X-12-ARIMA

Seasonal adjustment is done with X-12-ARIMA method.

Figure 3 WPI food articles (NSA and SA)




Figure 3 shows the non-seasonally and seasonally adjusted WPI Food. The plot reveals that the seasonal peaks are dampened after seasonal adjustment.

we also look at the plots of the growth rates to see the extent of noise reduction.

Figure 4 WPI (NSA and SA- point on point growth rates)




Figure 4 shows the point on point growth of the raw and seasonally adjusted series. The standard deviation of the growth rate of the raw series is 22.56 and that of the adjusted series is 11.62.


2.2 Diagnostic checks

After seasonal adjustment, a series of diagnostic checks are performed through relevant tests and quality assessment statistics.

2.2.1 Validation of the automodel choice by X-12-ARIMA

A test of validation of the auto model choice by X-12-ARIMA is the randomness of the residuals of the ARIMA model. The Ljung-Box test is conducted on the residuals of the fitted ARIMA model to check whether or not the residuals are white noise. The ACFs of the residuals are plotted to check for randomness.

Figure 5 ACF of residuals



The figure 5 does not reveal significant autocorrelation amongst the residuals.

2.2.2 Presence of identifiable seasonality

The statistic M7 shows the amount of moving seasonality present relative to stable seasonality. It shows the combined result for the test of stable and moving seasonality in the series. A value lesser than 0.7 is desirable to show identifiable seasonality in the series. The value of M7 is 0.342 for WPI Food

WPI Food series show identifiable seasonality on the basis of M7 statistic.

3 Year on year growth versus seasonally adjusted point on point growth

Growth rates can be computed either year on year or point on point. The year on year growth rate is computed as the percentage change with respect to the corresponding month (or quarter) in the preceding year, while the point on point growth rate is computed as the percentage change with respect to the preceding period.

Table 2 shows the year on year growth and seasonally adjusted annualized rate in percent, point on point.


4 Spectral representation

Figure 6 shows the spectral plot of the growth rate of the unadjusted and seasonally adjusted series. Spectral plot, an important tool of the frequency domain analysis shows the portion of variance of the series contributed by cycles of different frequencies.

The x-axis represent frequency from 0 to π (3.14). The seasonal frequencies are $\pi/6$ (0.52 on the x-axis), $\pi/3$ (1.04 on the x-axis), $\pi/2$ (1.57 on the x-axis), $2\pi/3$ (2.09 on the x-axis) and $5\pi/6$ (2.6 on the x-axis). In terms of periods (months); they are 12 months, 6 months, 4 months, 3 months and 2.4 months.

The figure at the lower panel shows that peaks at seasonal frequencies are eliminated after seasonal adjustment. For example the first peak at 0.52 correspond to 12 months which is eliminated after seasonal adjustment. Other peaks seen in the lower panel of the figure are not at seasonal frequencies.

Figure 6 WPI (Food) Spectral plot (NSA and SA)



5 Accounting for India-specific moving holiday effects

Accounting for moving holiday effect is a crucial component of pre-treatment of the series before the application of seasonal adjustment method. X-12-ARIMA is capable of handling the moving holiday effects through the inclusion of regressors for Easter Sunday, Labor Day,

and Thanksgiving Day. These are important moving holidays for U.S time series.

We use the GENHOL program of X-12-ARIMA to analyse India-specific moving holiday effect. The program generates regressor matrices from holiday date file to enable X-12-ARIMA, estimation of complex moving holiday effects. It has the capability to generate regressors for before the holiday interval, surrounding the holiday interval and past the holiday interval.

The key assumption is that the fundamental structure of a time series changes for a fixed number of days before, after or for a fixed interval surrounding the holidays. We estimate the effect of Diwali which is an important moving holiday in Indian scenario. We estimate the effect with different specifications about the number of days around the festival. However we did not find significant results for diwali effect on WPI (Food articles).

Table 1 Year on year and point on point growth rates

	Y.o.Y.growth	Point.on.point.growth
2006 Jan		9.62
2006 Feb		9.14
2006 Mar		-9.51
2006 Apr	5.98	-1.69
2006 May	7.98	19.88
2006 Jun	9.29	9.89
2006 Jul	5.35	8.46
2006 Aug	8.19	25.09
2006 Sep	10.68	15.28
2006 Oct	9.91	17.80
2006 Nov	9.10	5.13
2006 Dec	11.66	21.67
2007 Jan	11.43	8.26
2007 Feb	12.35	20.21
2007 Mar	13.24	0.11
2007 Apr	14.19	7.36
2007 May	12.93	5.52
2007 Jun	8.94	-4.53
2007 Jul	11.23	6.53
2007 Aug	9.75	9.49
2007 Sep	6.27	-4.51
2007 Oct	5.18	-13.32
2007 Nov	4.42	-5.15
2007 Dec	3.23	4.81
2008 Jan	0.84	14.35
2008 Feb	2.77	13.99
2008 Mar	5.63	32.22
2008 Apr	6.09	11.04
2008 May	6.46	9.98
2008 Jun	6.98	-0.01
2008 Jul	6.89	7.28
2008 Aug	6.58	7.28
2008 Sep	8.21	12.74
2008 Oct	11.35	22.28
2008 Nov	12.70	7.56
2008 Dec	12.09	-5.14
2009 Jan	14.43	9.68
2009 Feb	9.47	-4.76
2009 Mar	7.96	14.08
2009 Apr	8.69	18.73
2009 May	8.91	13.12
2009 Jun	11.28	23.51
2009 Jul	12.74	24.96
2009 Aug	14.36	26.83
2009 Sep	13.92	5.84
2009 Oct	12.47	32.35
2009 Nov	16.73	26.17
2009 Dec	20.76	32.93
2010 Jan	20.19	5.88
2010 Feb	21.85	13.96
2010 Mar	20.65	1.02
2010 Apr	20.49	16.60